

Analysis of 62-day pathways using 2017-2018 Cancer Waiting Times data for colorectal, lung and prostate cancers

England, by Cancer Alliance

Produced by the Cancer Alliance Data, Evidence and Analysis Service

19 December, 2018



Public Health
England



Contents

- ▶ Background
- ▶ Methodology
- ▶ Summary of key findings
- ▶ Count of patients with a CWT record included in the analysis, by Cancer Alliance and cancer type

The following graphs are presented for colorectal, lung and prostate cancers separately:

- ▶ Variation in days from a referral to first treatment received, by Cancer Alliance
- ▶ Variation in first treatment modality received, by Cancer Alliance
- ▶ Variation in days from a referral to first treatment received, by five most common treatment modalities
- ▶ Variation in days from a referral to first treatment received, by Cancer Alliance, for the most common treatment modality
- ▶ Variation in days from a referral to first treatment received, by Cancer Alliance, for the second most common treatment modality
- ▶ Variation in days from a referral to first treatment received, by Cancer Alliance, for the third most common treatment modality

Background

Aims: to provide Cancer Alliances with analysis of pathways with a 62-day¹ Cancer Waiting Times (CWT) record for colorectal, lung and prostate cancers. This is to enable Alliances to support:

- ▶ implementation of the best practice times pathways; and
- ▶ meeting 62-day CWT operational standards.

The work is also aimed at helping Cancer Alliances understand where variations are occurring across Alliances, as well as understanding pathway variation between cancer types.

Cancer Alliances are encouraged to use the analysis to identify best practice and to investigate significant variations where appropriate.

This work is produced by the Cancer Alliance Data, Evidence and Analysis Service (CADEAS), a partnership between NHS England and Public Health England. It builds on work previously carried out by RM Partners for the three Cancer Vanguard sites.

¹The 62 day CWT target states that 85% of patients who are urgently referred by their GP for suspected cancer must have their first definitive treatment within 62 days.

This analysis uses CWT dataset records for patients diagnosed with colorectal (C18-C20), lung (C34) or prostate (C61) cancers². Only patients on a 62-day pathway were included in this analysis³.

Only records with a first treatment date recorded between 1st April 2017 and 31st March 2018 were included.

The work is segmented by:

- ▶ Cancer type
- ▶ Cancer Alliance
- ▶ Days from referral to first treatment
- ▶ First treatment modality received

Two CWT adjustments have been included in the time taken from referral to first treatment calculations, in line with CWT official statistics⁴.

²Please note that a small proportion of patients (0.1%) have multiple CWT records; this is due to patients being diagnosed with more than one tumour or having multiple CWT data submissions for the same tumour.

³Please note that patients with a CWT record account for approximately 80% of those diagnosed with cancer in England.

⁴National Cancer Waiting Times Monitoring Dataset Guidance - Version 9.0 (pg.47)

Methodology (cont.)

Cancer Alliances will receive a national report for England segmented by Cancer Alliance and three local reports segmented by:

- ▶ Resident clinical commissioning group (CCG)
- ▶ Treatment trust
- ▶ Trust first seen

This report should be read in conjunction with the guidance document.

Figures for Cancer Alliances as a whole are derived from their commissioner organisations. The England figure shown is the total for all patients treated by English trusts (this includes those patients with an unknown commissioner) as this is the official position for England performance. Patients registered to Welsh commissioners or private trusts were excluded from the analysis. Data completeness in the cohort for England was 98.44% in the CCG field, 100% in the treatment trust field and 99.92% in the trust first seen field.

Caveats:

- ▶ This is operational data to be used for operational decision making and for this reason has not been casemix adjusted.
- ▶ Caution should be taken when interpreting results with small cohorts as small numbers can lead to variation and unreliability of data. In cases where there are five or fewer patients, the patient number is recorded as <6.


Summary of key findings

A total of 47948 patients were included in the analysis for 2017/18 (28% colorectal; 23% lung; and 49% prostate cancers). East of England Cancer Alliance had the highest number of cancer cases with a CWT record in 2017/18 for all three cancer sites (colorectal, lung and prostate cancers) (slide 7).

The three most common first treatment modalities received for colorectal cancers are surgery, palliative care only and chemotherapy; chemotherapy, surgery and palliative care only for lung cancer; and hormonal therapy, active monitoring and surgery for prostate cancers (slides 12-14, 19-21 and 26-28).


There is variation in days from a referral to first treatment received across Cancer Alliances even for the same cancer type and same treatment type. This may be due to stage at diagnosis and patient characteristics. It is important to note that a shorter pathway may not always be associated with better outcomes. Further analysis will be carried out to examine variation by stage at diagnosis and patient characteristics using linked datasets.

Count of patients with a CWT record included in the analysis, by Cancer Alliance and cancer type (2017/18)




COLORECTAL CANCER

Colorectal cancer: variation in days from a referral to first treatment received, by Cancer Alliance (2017/18)




Figures within the bars represent the median values for each group.


Colorectal cancer: variation in first treatment modality received, by Cancer Alliance (2017/18)



Colorectal cancer: variation in days from a referral to first treatment received, by the five most common treatment modalities (2017/18)




Colorectal cancer and surgery: variation in days from a referral to first treatment received, by Cancer Alliance (2017/18)




Figures within the bars represent the median values for each group.

Colorectal cancer and palliative care only: variation in days from a referral to first treatment received, by Cancer Alliance (2017/18)



Figures within the bars represent the median values for each group.


Colorectal cancer and chemotherapy: variation in days from a referral to first treatment received, by Cancer Alliance (2017/18)



Figures within the bars represent the median values for each group.


LUNG CANCER

Lung cancer: variation in days from a referral to first treatment received, by Cancer Alliance (2017/18)




Figures within the bars represent the median values for each group.

Lung cancer: variation in first treatment modality received, by Cancer Alliance (2017/18)




Lung cancer: variation in days from a referral to first treatment received, by the five most common treatment modalities (2017/18)




Figures within the bars represent the median values for each group.

Lung cancer and chemotherapy: variation in days from a referral to first treatment received, by Cancer Alliance (2017/18)




Figures within the bars represent the median values for each group.

Lung cancer and surgery: variation in days from a referral to first treatment received, by Cancer Alliance (2017/18)



Figures within the bars represent the median values for each group.


Lung cancer and palliative care only: variation in days from a referral to first treatment received, by Cancer Alliance (2017/18)



Figures within the bars represent the median values for each group.


PROSTATE CANCER

Prostate cancer: variation in days from a referral to first treatment received, by Cancer Alliance (2017/18)




Figures within the bars represent the median values for each group.

Prostate cancer: variation in first treatment modality received, by Cancer Alliance (2017/18)




Prostate cancer: variation in days from a referral to first treatment received, by the five most common treatment modalities (2017/18)




Figures within the bars represent the median values for each group.

Prostate cancer and hormonal therapy: variation in days from a referral to first treatment received, by Cancer Alliance (2017/18)




Figures within the bars represent the median values for each group.

Prostate cancer and active monitoring: variation in days from a referral to first treatment received, by Cancer Alliance (2017/18)



Figures within the bars represent the median values for each group.

Prostate cancer and surgery: variation in days from a referral to first treatment received, by Cancer Alliance (2017/18)



Figures within the bars represent the median values for each group.